arXiv:0903.3151v1 [math.CO] 18 Mar 2009

ON THE VERTEX FOLKMAN NUMBERS $F_v(\underbrace{2,\dots,2}_{r};r-1)$ and $F_v(\underbrace{2,\dots,2}_{r};r-2)$*

Nedyalko Nenov

March 18, 2009

Abstract

For a graph G the symbol $G \overset{v}{\to} (a_1,\dots,a_r)$ means that in every r-coloring of the vertices of G, for some $i \in \{1,2,\dots,r\}$, there exists a monochromatic a_i-clique of color i. The vertex Folkman numbers

$$F_v(a_1,\dots,a_r;q) = \min\{|V(G)| : G \overset{v}{\to} (a_1,\dots,a_r) \text{ and } K_q \nsubseteq G\}$$

are considered. We prove that $F_v(\underbrace{2,\dots,2}_{r};r-1) = r+7$, $r \geq 6$ and $F_v(\underbrace{2,\dots,2}_{r};r-2) = r+9$, $r \geq 8$.

Keywords. Folkman graphs, Folkman numbers
2000 Math. Subject Classification. 05C55

1 Introduction

We consider only finite, non-oriented graphs without loops and multiple edges. We call a p-clique of the graph G a set of p vertices, each two of which are adjacent. The largest positive integer p, such that the graph G contains a p-clique is denoted by $\mathrm{cl}(G)$. In this paper we shall also use the following notations:

- $V(G)$ is the vertex set of the graph G;
- $E(G)$ is the edge set of the graph G;
- $\overline{G}$ is the complement of G;
- $G[V]$, $V \subseteq V(G)$ is the subgraph of G induced by V;

---

*This work is supported by the Scientific Research Fund of the St. Kl. Ohridski Sofia University under contracts 85/2006.

- $G - V$, $V \subseteq V(G)$ is the subgraph of G induced by $V(G) \setminus V$;
- $\alpha(G)$ is the vertex independence number of G;
- $\chi(G)$ is the chromatic number of G;
- $f(G) = \chi(G) - \mathrm{cl}(G)$;
- K_n is the complete graph on n vertices;
- C_n is the simple cycle on n vertices.

Let G_1 and G_2 be two graphs without common vertices. We denote by $G_1 + G_2$ the graph G for which $V(G) = V(G_1) \cup V(G_2)$ and $E(G) = E(G_1) \cup E(G_2) \cup E'$, where $E' = \{[x, y] : x \in V(G_1), y \in V(G_2)\}$.

The Ramsey number $R(p, q)$ is the smallest natural n such that for every n-vertex graph G either $\mathrm{cl}(G) \geq p$ or $\alpha(G) \geq q$. An exposition of the results on the Ramsey numbers is given in [25]. We shall need Table 1.1 of the known Ramsey numbers $R(p, 3)$ (see [25]).

p	3	4	5	6	7	8	9	10
$R(p,3)$	6	9	14	18	23	28	36	40–43

Table 1.1: Known Ramsey numbers

Definition. Let $a_1, \dots, a_r$ be positive integers. We say that the r-coloring

$$V(G) = V_1 \cup \dots \cup V_r, \quad V_i \cap V_j = \emptyset,\ i \neq j$$

of the vertices of the graph G is $(a_1, \dots, a_r)$-free, if V_i does not contain an a_i-clique for each $i \in \{1, \dots, r\}$. The symbol $G \xrightarrow{v} (a_1, \dots, a_r)$ means that there is no $(a_1, \dots, a_r)$-free coloring of the vertices of G.

Let $a_1, \dots, a_r$ and q be natural numbers. Define

$$H_v(a_1, \dots, a_r; q) = \{G : G \xrightarrow{v} (a_1, \dots, a_r) \text{ and } \mathrm{cl}(G) < q\},$$
$$F_v(a_1, \dots, a_r; q) = \min\{|V(G)| : G \in H_v(a_1, \dots, a_r; q)\}.$$

We say that the graph $G \in H_v(a_1, \dots, a_r; q)$ is an extremal graph in $H_v(a_1, \dots, a_r; q)$, if $|V(G)| = F_v(a_1, \dots, a_r; q)$.

It is clear that $G \xrightarrow{v} (a_1, \dots, a_r)$ implies $\mathrm{cl}(G) \geq \max\{a_1, \dots, a_r\}$. Folkman [3] proved that there exists a graph G such that $G \xrightarrow{v} (a_1, \dots, a_r)$ and $\mathrm{cl}(G) = \max\{a_1, \dots, a_r\}$. Therefore

$$F_v(a_1, \dots, a_r; q) \text{ exists} \iff q > \max\{a_1, \dots, a_r\}. \tag{1.1}$$

The numbers $F_v(a_1, \dots, a_r; q)$ are called vertex Folkman numbers.

If $a_1, \dots, a_r$ are positive integers, $r \geq 2$ and $a_i = 1$ then it is easily to see that

$$G \xrightarrow{v} (a_1, \dots, a_i, \dots, a_r) \iff G \xrightarrow{v} (a_1, \dots, a_{i-1}, a_{i+1}, a_r).$$

Thus it is enough to consider just such numbers $F_v(a_1, \dots, a_r; q)$ for which $a_i \geq 2$, $i = 1, \dots, r$. In this paper we consider the vertex Folkman numbers $F_v(2, \dots, 2; q)$. Set

$$(\underbrace{2, \dots, 2}_{r}) = (2_r) \quad \text{and} \quad F_v(\underbrace{2, \dots, 2}_{r}; q) = F_v(2_r; q).$$

By (1.1)

$$F_v(2_r; q) \text{ exists} \iff q \geq 3. \tag{1.2}$$

It is clear that

$$G \xrightarrow{v} (2_r) \iff \chi(G) \geq r + 1. \tag{1.3}$$

Since $K_{r+1} \xrightarrow{v} (2_r)$ and $K_r \overset{v}{\not\rightarrow} (2_r)$ we have

$$F_v(2_r; q) = r + 1 \quad \text{if} \quad q \geq r + 2.$$

In [2] Dirac proved the following

Theorem 1.1 ([2])**.** *Let G be a graph such that $\chi(G) \geq r+1$ and $\mathrm{cl}(G) \leq r$. Then*

(a) $|V(G)| \geq r + 3$*;*

(b) *If $|V(G)| = r + 3$ then $G = K_{r-3} + C_5$.*

According to (1.3), Theorem 1.1 is equivalent to the following

Theorem 1.2. *Let $r \geq 2$ be a positive integer. Then*

(a) $F_v(2_r; r + 1) = r + 3$*;*

(b) *$K_{r-3} + C_5$ is the only extremal graph in $H_v(2_r; r + 1)$.*

In [14] Łuczak, Ruciński and Urbański defined for arbitrary positive integers $a_1, \dots, a_r$ the numbers

$$m = \sum_{i=1}^{r}(a_i - 1) + 1 \quad \text{and} \quad p = \max\{a_1, \dots, a_r\}. \tag{1.4}$$

and proved the following extension of Theorem 1.2.

Theorem 1.3 ([14])**.** *Let $a_1, \dots, a_r$ be positive integers and m and p be defined by* (1.4)*. Let $m \geq p + 1$. Then*

(a) $F_v(a_1, \dots, a_r; m) = m + p$*;*

(b) *$K_{m-p-1} + \overline{C}_{2p+1}$ is the only extremal graph in $H_v(a_1, \dots, a_r; m)$.*

An another extension of Theorem 1.1 was given in [21].

From (1.1) it follows that the numbers $F_v(a_1, \dots, a_r; m - 1)$ exists if and only if $m \geq p + 2$. The exact values of all numbers $F_v(a_1, \dots, a_r; m - 1)$ for which $p = \max\{a_1, \dots, a_r\} \leq 4$ are known. A detailed exposition of these results was given in [13] and [23]. We do not know any exact values of $F_v(a_1, \dots, a_r; m-1)$ in the case when $\max\{a_1, \dots, a_r\} \geq 5$. Here we shall note only the values $F_v(a_1, \dots, a_r; m-1)$ when $a_1 = a_2 = \cdots = a_r = 2$, i.e. of the numbers $F_v(2_r; r)$. From (1.2) these numbers exist if and only if $r \geq 3$. If $r = 3$ and $r = 4$ we have that

$$F_v(2_3; 3) = 11; \tag{1.5}$$

$$F_v(2_4; 4) = 11. \tag{1.6}$$

The inequality $F_v(2_3; 3) \leq 11$ was proved in [15] and the opposite inequality $F_v(2_3; 3) \geq 11$ was proved in [1]. The equality (1.6) was proved in[18] (see also [19]). If $r \geq 5$ we have the following

Theorem 1.4 ([17], see also [24])**.** *Let* $r \geq 5$*. Then:*

(a) $F_v(2_r; r) = r + 5$*;*

(b) $K_{r-5} + C_5 + C_5$ *is the only extremal graph in* $H_v(2_r; r)$*.*

Theorem 1.4(a) was proved also in [8] and [14].

According to (1.2) the number $F_v(2_r; r-1)$ exists if and only if $r \geq 4$. In [17] we prove that

$$F_v(2_r; r-1) = r + 7 \text{ if } r \geq 8. \tag{1.7}$$

In this paper we shall improve (1.7) proving the following

Theorem 1.5. *Let* $r \geq 4$ *be an integer. Then*

(a) $F_v(2_r; r-1) \geq r + 7$*;*

(b) $F_v(2_r; r-1) = r + 7$ *if* $r \geq 6$*;*

(c) $F_v(2_5; 4) \leq 16$.

In [9] Jensen and Royle show that

$$F_v(2_4; 3) = 22. \tag{1.8}$$

We see from Theorem 1.5 and (1.8) that $F_v(2_5; 4)$ is the only unknown number of the kind $F(2_r; r-1)$.

From (1.2) the Folkman number $F(2_r; r-2)$ exists if and only if $r \geq 5$. In [16] we prove that $F_v(2_r; r-2) = r + 9$ if $r \geq 11$. In this paper we shall improve this result proving the folloing

Theorem 1.6. *Let* $r \geq 5$ *be an integer. Then*

(a) $F_v(2_r; r-2) \geq r + 9$*;*

(b) $F_v(2_r; r-2) = r + 9$ *if* $r \geq 8$.

The numbers $F_v(2_r; r-2)$, $5 \leq r \leq 7$, are unknown.

2 Auxiliary results

Let G be an arbitrary graph. Define

$$f(G) = \chi(G) - \mathrm{cl}(G).$$

Lemma 2.1. *Let G be a graph such that $f(G) \leq 2$. Then $|V(G)| \geq \chi(G) + 2f(G)$.*

Proof. Since $\chi(G) \geq \mathrm{cl}(G)$, $f(G) \geq 0$. If $f(G) = 0$ the inequality is trivial. Let $f(G) = 1$ and $\chi(G) = r + 1$. Then $\mathrm{cl}(G) = r$. Note that $r \geq 2$ because $\chi(G) \neq \mathrm{cl}(G)$. By (1.3), $G \in H_v(2_r; r + 1)$. Thus, from Theorem 1.2(a), it follows $|V(G)| \geq r+3 = 2f(G)+\chi(G)$. Let $f(G) = 2$ and $\chi(G) = r+1$. Then $\mathrm{cl}(G) = r-1$. Since $\chi(G) \neq \mathrm{cl}(G)$, $\mathrm{cl}(G) = r - 1 \geq 2$, i.e. $r \geq 3$. From Theorem 1.4(a), (1.5) and (1.6) we obtain that $|V(G)| \geq r + 5 = \chi(G) + 2f(G)$. This completes the proof of Lemma 2.1. □

Let $G = G_1 + G_2$. Obviously,

$$\chi(G) = \chi(G_1) + \chi(G_2); \tag{2.1}$$

$$\mathrm{cl}(G) = \mathrm{cl}(G_1) + \mathrm{cl}(G_2). \tag{2.2}$$

Hence,

$$f(G) = f(G_1) + f(G_2). \tag{2.3}$$

A graph G is defined to be vertex-critical chromatic if $\chi(G - v) < \chi(G)$ for all $v \in V(G)$. We shall use the following result in the proof of Theorem 1.6.

Theorem 2.1 ([4], see also [5])**.** *Let G be a vertex-critical chromatic graph and $\chi(G) \geq 2$. If $|V(G)| < 2\chi(G) - 1$, then $G = G_1 + G_2$, where $V(G_i) \neq \emptyset$, $i = 1, 2$.*

Remark. In the original statement of Theorem 2.1 the graph G is edge-critical chromatic (and not vertex-critical chromatic). Since each vertex-critical chromatic graph G contains an edge-critical chromatic subgraph H such that $\chi(G) = \chi(H)$ and $V(G) = V(H)$, the above statement of this theorem is equivalent to the original one. They are also more convenient for the proof of Theorem 1.6.

Let G be a graph and $A \subseteq V(G)$ be an independent set of vertices of the graph G. It is easy to see that

$$G \xrightarrow{v} (2_r),\ r \geq 2 \Rightarrow G - A \xrightarrow{v} (2_{r-1}). \tag{2.4}$$

Lemma 2.2. *Let $G \in H_v(2_r; q)$, $q \geq 3$ and $|V(G)| = F_v(2_r; q)$. Then*

(a) *G is a vertex-critical $(r + 1)$-chromatic graph;*

(b) *if $q < r + 3$ then $\mathrm{cl}(G) = q - 1$.*

Proof. By (1.3), $\chi(G) \geq r + 1$. Assume that (a) is wrong. Then there exists $v \in V(G)$ such that $\chi(G - v) \geq r + 1$. According to (1.3), $G - v \in H_v(2_r; q)$. This contradicts the equality $|V(G)| = F_v(2_r; q)$.

Assume that (b) is wrong, i.e. $\mathrm{cl}(G) \leq q-2$. Then from $q < r+3$ it follows that $\mathrm{cl}(G) < r+1$. Since $\chi(G) \geq r+1$ there are $a, b \in V(G)$ such that $[a,b] \notin E(G)$. Consider the subgraph $G_1 = G - \{a,b\}$. We have $r \geq 2$, because $\chi(G) \neq \mathrm{cl}(G)$. Thus, from (2.4) and $\mathrm{cl}(G) \leq q-2$ it follows that $G_1 \in H_v(2_{r-1}; q-1)$. Obviously, $G_1 \in H_v(2_{r-1}; q-1)$ leads to $K_1 + G_1 \in H_v(2_r; q)$. This contradicts the equality $|V(G)| = F_v(2_r; q)$, because $|V(K_1 + G_1)| = |V(G)| - 1$. Lemma 2.2 is proved. $\square$

Lemma 2.3. *Let* $G \in H_v(2_r; q)$, $r \geq 2$. *Then*

$$|V(G)| \geq F_v(2_{r-1}; q) + \alpha(G).$$

Proof. Let $A \subseteq V(G)$ be an independent set such that $|A| = \alpha(G)$. Consider the subgraph $G_1 = G - A$. According to (2.4), $G_1 \in H_v(2_{r-1}; q)$. Hence $|V(G_1)| \geq F_v(2_{r-1}; q)$. Since $|V(G)| = |V(G_1)| + \alpha(G)$, Lemma 2.3 is proved. $\square$

We shall use also the following three results:

$$F_v(2,2,p;p+1) \geq 2p+4, \qquad \text{see [20];} \tag{2.5}$$

$$F_v(2,2,4;5) = 13, \qquad \text{see [22].} \tag{2.6}$$

Theorem 2.2 ([12]). *Let* G *be a graph,* $\mathrm{cl}(G) \leq p$ *and* $|V(G)| \geq p+2$, $p \geq 2$. *Let* G *also have the following two properties:*

(i) $G \overset{v}{\not\to} (2,2,p)$;

(ii) *If* $V(G) = V_1 \cup V_2 \cup V_3$ *is a* $(2,2,p)$*-free 3-coloring then* $|V_1| + |V_2| \leq 3$.

Then $G = K_1 + G_1$.

3 An upper bound for the numbers $F_v(2_r; q)$

Consider the graph P, whose complementary graph $\overline{P}$ is given in Fig. 1. This



Figure 1: Graph $\overline{P}$

graph is a well-known construction of Greenwood and Gleason [6], which shows that $R(5,3) \geq 14$, because $|V(P)| = 13$ and

$$\alpha(P) = 2; \tag{3.1}$$

$$\mathrm{cl}(P) = 4 \qquad \text{see [6].} \tag{3.2}$$

From $|V(P)| = 13$ and (3.1) it follows that $\chi(P) \geq 7$. Since $\{v_1\} \cup \{v_2, v_3\} \cup \cdots \cup \{v_{12}, v_{13}\}$ is a 7-chromatic partition of $V(P)$, we have

$$\chi(P) = 7. \tag{3.3}$$

Let r and s be non-negative integers and $r \geq 3s + 6$. Define

$$\tilde{P} = K_{r-3s-6} + P + \underbrace{C_5 + \cdots + C_5}_{s}.$$

From (2.1), (2.2), (3.2) and (3.3) we obtain that $\chi(\tilde{P}) = r+1$ and $\mathrm{cl}(\tilde{P}) = r-s-2$. By (1.3) $\tilde{P} \in H_v(2_r; r-s-1)$ and thus

$$F_v(2_r; r-s-1) \leq |V(\tilde{P})|.$$

Since $|V(\tilde{P})| = r + 2s + 7$, we prove the following

Theorem 3.1. *Let r and s be non-negative integers and $r \geq 3s + 6$. Then*

$$F_v(2_r; r-s-1) \leq r + 2s + 7.$$

Remark. Since $r \geq 3s+6$ we have $r - s - 1 > 2$. Thus, according to (1.2), the numbers $F_v(2_r; r-s-1)$ exists.

4 Proof of Theorem 1.5

Proof of Theorem 1.5(a). Let $G \in H_v(2_r; r-1)$. We need to prove that $|V(G)| \geq r + 7$. From Lemma 2.3 we have that

$$|V(G)| \geq F_v(2_{r-1}; r-1) + \alpha(G).$$

By (1.5), (1.6) and Theorem 1.4(a), $F_v(2_{r-1}; r-1) \geq r+4$. Hence

$$|V(G)| \geq r + 4 + \alpha(G). \tag{4.1}$$

We prove the inequality $|V(G)| \geq r+7$ by induction on r. From Table 1.1 we see that

$$R(r-1,3) < r+6 \text{ if } r=4 \text{ or } r=5. \tag{4.2}$$

Obviously, from $G \in H_v(2_r; r-1)$ it follows that $\chi(G) \neq \mathrm{cl}(G)$. Thus, $\alpha(G) \geq 2$. From (4.1) we obtain $|V(G)| \geq r+6$. From this inequality and (4.2) we see that

$|V(G)| > R(r-1, 3)$ if $r = 4$ or $r = 5$. Since $\mathrm{cl}(G) < r-1$, it follows that $\alpha(G) \geq 3$. Now from (4.1) we obtain that $|V(G)| \geq r + 7$ if $r = 4$ or $r = 5$.

Let $r \geq 6$. We shall consider two cases:

Case 1. $G \overset{v}{\not\to} (2, 2, r-2)$. From Theorem 2.2 we see that only following two subcases are possible:

Subcase 1a. $G = K_1 + G_1$. From $G \in H_v(2_r, r-1)$ it follows that $G_1 \in H_v(2_{r-1}; r-2)$. By the inductive hypothesis, $|V(G_1)| \geq r+6$. Therefore, $|V(G)| \geq r + 7$.

Subcase 1b. There is a $(2, 2, r-2)$-free 3-coloring $V(G) = V_1 \cup V_2 \cup V_3$ such that $|V_1| + |V_2| \geq 4$. Consider the subgraph $\tilde{G} = G[V_3]$. By assumption $\tilde{G}$ does not contain an $(r-2)$-clique, i.e. $\mathrm{cl}(\tilde{G}) < r-2$. Since V_1 and V_2 are independent sets and $G \xrightarrow{v} (2_r)$ it follows from (2.4) that $\tilde{G} \xrightarrow{v} (2_{r-2})$. Thus, $\tilde{G} \in H_v(2_{r-2}; r-2)$. By (1.6) and Theorem 1.4(a), $|V(\tilde{G})| \geq r+3$. As $|V_1|+|V_2| \geq 4$, we have $|V(G)| \geq 7$.

Case 2. $G \xrightarrow{v} (2, 2, r-2)$. Since $\mathrm{cl}(G) < r-1$, $G \in H_v(2, 2, r-2; r-1)$. From (2.5) it follows that $|V(G)| \geq 2(r-2) + 4 = 2r$. Hence, if $2r \geq r + 7$, i.e. $r \geq 7$, we have $|V(G)| \geq r + 7$. Let $r = 6$. Then $G \in H_v(2, 2, 4; 5)$. By (2.6), $|V(G)| \geq 13$. □

Proof of Theorem 1.5(b). Let $r \geq 6$. According to Theorem 1.5(a) we have $F_v(2_r; r-1) \geq r + 7$. From Theorem 3.1 ($s = 0$) we obtain the opposite inequality $F_v(2_r; r-1) \leq r + 7$. □

Proof of Theorem 1.5(c). There is a 16-vertex graph G such that $\alpha(G) = 3$ and $\mathrm{cl}(G) = 3$, because $R(4, 4) = 18$ (see [6]). From $|V(G)| = 16$ and $\alpha(G) = 3$ obviously it follows that $\chi(G) \geq 6$. By (1.3), $G \xrightarrow{v} (2_5)$. So, $G \in H_v(2_5; 4)$. Hence $F_v(2_5; 4) \leq |V(G)| = 16$. □

Theorem 1.5 is proved.

Corollary 4.1. *Let G be a graph such that $f(G) \leq 3$. Then $|V(G)| \geq \chi(G)+2f(G)$.*

Proof. If $f(G) \leq 2$, then Corollary 4.1 follows from Lemma 2.1. Let $f(G) = 3$ and $\chi(G) = r + 1$. Then $\mathrm{cl}(G) = r - 2$. Since $\chi(G) \neq \mathrm{cl}(G)$, $\mathrm{cl}(G) \geq 2$. Thus, $r \geq 4$. By (1.3), $G \in H_v(2_r; r-1)$. From Theorem 1.5(a) we obtain $|V(G)| \geq r + 7 = \chi(G) + 2f(G)$. □

Remark. In $H_v(2_r; r-1)$, $r \geq 8$ there are more than one extremal graph. For example, in $H_v(2_8; 7)$ besides $K_2 + P$ (see Theorem 3.1), the graph $C_5 + C_5 + C_5$ is extremal, too.

5 Proof of Theorem 1.6

Proof of Theorem 1.6(a). Let $G \in H_v(2_r; r-2)$. We need to prove that $|V(G)| \geq r + 9$. From Lemma 2.3 we have that

$$|V(G)| \geq F_v(2_{r-1}; r-2) + \alpha(G).$$

By Theorem 1.5(a), $F_v(2_{r-1}; r-2) \geq r+6$. Thus,

$$|V(G)| \geq r + 6 + \alpha(G). \tag{5.1}$$

We prove the inequality $|V(G)| \geq r+9$ by induction on r. From Table 1.1 we see that

$$R(r-2,3) < r+8, \quad 5 \leq r \leq 7. \tag{5.2}$$

Obviously, from $G \in H_v(2_r; r-2)$ it follows that $\chi(G) \neq \mathrm{cl}(G)$. Thus, $\alpha(G) \geq 2$. From (5.1) we obtain $|V(G)| \geq r+8$. This, together with (5.2), implies $|V(G)| > R(r-2,3)$ if $5 \leq r \leq 7$. Since $\mathrm{cl}(G) < r-2$, $\alpha(G) \geq 3$. By the inequality (5.1), $|V(G)| \geq r+9$, $5 \leq r \leq 7$.

Let $r \geq 8$. Obviously, it is enough to consider only the situation when

$$|V(G)| = F_v(2_r; r-2). \tag{5.3}$$

By (5.3) and Lemma 2.2 we have that

$$G \textit{ is a vertex-critical } (r+1)\textit{-chromatic graph;} \tag{5.4}$$

and

$$\mathrm{cl}(G) = r-3. \tag{5.5}$$

From (5.4) and (5.5) it follows that

$$f(G) = 4. \tag{5.6}$$

We shall consider two cases.

Case 1. $|V(G)| < 2r+1$. By (5.4) and Theorem 2.1, we obtain that

$$G = G_1 + G_2. \tag{5.7}$$

From (5.7), (2.1) and (5.4), obviously it follows that

$$G_i,\ i=1,2 \textit{ is a vertex-critical chromatic graph.} \tag{5.8}$$

Let $f(G_1) = 0$. Then, according to (5.8) G_1 is a complete graph. Thus, it follows from (5.7) that $G = K_1 + G'$. It is clear that

$$G \in H_v(2_r; r-2) \Rightarrow G' \in H_v(2_{r-1}; r-3).$$

By the inductive hypothesis, $|V(G')| \geq r+8$. Hence, $|V(G)| \geq r+9$. Let $f(G_i) \neq 0$, $i=1,2$. We see from (5.7), (2.3) and (5.6) that $f(G_i) \leq 3$, $i=1,2$. By Corollary 4.1, we have

$$|V(G_i)| \geq \chi(G_i) + 2f(G_i), \quad i=1,2.$$

Summing these inequalities and using (2.1) and (2.3) we obtain

$$|V(G)| \geq \chi(G) + 2f(G). \tag{5.9}$$

According to (5.4), $\chi(G) = r+1$. Thus, from (5.9) and (5.6) it follows that $|V(G)| \geq r+9$.

Case 2. $|V(G)| \geq 2r+1$. Since $r \geq 8$, $2r+1 \geq r+9$. Hence $|V(G)| \geq r+9$. □

Proof of Theorem 1.6(b). By Theorem 1.6(a), $F_v(2_r; r-2) \geq r+9$. Thus, we need to prove the opposite inequality $F_v(2_r; r-2) \leq r+9$ if $r \geq 8$. If $r \geq 9$, this inequality follows from Theorem 3.1 ($s=1$). Let $r=8$. By $R(6,3)=18$, [11] (see also [7]), there is a graph Q such that $|V(Q)|=17$, $\alpha(Q)=2$ and $\mathrm{cl}(Q)=5$. From $|V(Q)|=17$ and $\alpha(Q)=2$ obviously it follows that $\chi(Q) \geq 9$. Thus, by (1.3), $Q \xrightarrow{v} (2_8)$. Hence $Q \in H_v(2_8; 6)$ and $F_v(2_8; 6) \leq |V(Q)| = 17$. Theorem 1.6 is proved. □

Corollary 5.1. *Let G be a graph such that $f(G) \leq 4$. Then $|V(G)| \geq \chi(G)+2f(G)$.*

Proof. If $f(G) \leq 3$, then Corollary 5.1 follows from Corollary 4.1. Let $f(G)=4$ and $\chi(G)=r+1$. Then $\mathrm{cl}(G)=r-3$. Since $\chi(G) \neq \mathrm{cl}(G)$, $\mathrm{cl}(G) \geq 2$. Thus, $r \geq 5$. By (1.3), $G \in H_v(2_r; r-2)$. Using Theorem 1.6(a), we get $|V(G)| \geq r+9 = \chi(G)+2f(G)$. □

Let $r \geq 3s+8$. Define

$$\tilde{Q} = K_{r-3s-8} + Q + \underbrace{C_5 + \cdots + C_5}_{s},$$

where graph Q is given in the proof of Theorem 1.6(b). Since $\mathrm{cl}(Q)=5$ and $\chi(Q) \geq 9$, we have by (2.1) and (2.2) that $\mathrm{cl}(\tilde{Q}) = r-s-3$ and $\chi(\tilde{Q}) \geq r+1$. According to (1.3), $\tilde{Q} \in H_v(2_r; r-s-2)$. Thus, $F_v(2_r; r-s-2) \leq |V(\tilde{Q})|$. Since $|V(\tilde{Q})| = r+2s+9$, we obtain the following

Theorem 5.1. *Let r and s be non-negative integers and $r \geq 3s+8$. Then*

$$F_v(2_r; r-s-2) \leq r+2s+9.$$

References

[1] Chvátal, V. The minimality of the Mycielski graph. *Lecture Notes Math.*, **406**, 1979, 243–246.

[2] Dirac, G. Map colour theorems related to the Heawood colour formula. *J. London Math. Soc.*, **31**, 1956, 460–471.

[3] Folkman, J. Graphs with monochromatic complete subgraphs in every edge coloring. *SIAM J. Appl. Math.* **18**, 1970, 19–24.

[4] Gallai, T. Kritische graphen II, *Publ. Math. Inst. Hungar. Acad. Sci.*, **8**, 1963, 373–395.

[5] Gallai, T. Critical Graphs. In: Theory of Graphs and Its Applications, Proceedings of the Symposium held in Smolenice in June 1963, Czechoslovak Acad. Sciences, Prague, 1964, pp. 43–45.

[6] Greenwood, R., A. Gleason. Combinatorial relation and chromatic graphs. *Canad. J. Math.*, **7**, 1955, 1–7.

[7] Grinstead, C., S. Roberts. On the Ramsey numbers $R(3, 8)$ and $R(3, 9)$. *J. Comb. Theory*, **B33**, 1982, 27–51.

[8] Guta, P. On the structure of k-chromatic critical graphs of order $k + p$. *Stud. Cerc. Math.*, **50**, 1998, No 3–4, 169–173.

[9] Jensen, T., G. Royle. Small graphs with chromatic number 5: a computer research. *J. Graph Theory*, **19**, 1995, 107–116.

[10] Kalbfleish, J. On an Unknown Ramsey Number. *Michigan Math. J.*, **13**, 1966, 385–392.

[11] Kery, G. On a theorem of Ramsey. *Mat. Lapok*, **15**, 1964, 204–224.

[12] Kolev, N., N. Nenov. Folkman number $F_e(3, 4; 8)$ is equal to 16. *Compt. rend. Acad. bulg. Sci.*, **59**, 2006, No 1, 25–30.

[13] Kolev, N., N. Nenov. New recurrent Inequality on a Class of Vertex Folkman Numbers. In: Proc. 35th Spring Conf. of Union Bulg. Mathematicians, Borovets, April 5–8, 2006, pp. 164–168.

[14] Łuczak, T., A. Ruciński, S. Urbański. On minimal vertex Folkman graphs. *Discrete Math.*, **236**, 2001, 245–262.

[15] Mycielski, J. Sur le coloriage des graphes. *Colloq. Math.*, **3**, 1955, 161–162.

[16] Nenov, N. Lower bound for some constants related to Ramsey graphs. *Annuaire Univ. Sofia Fac. Math. Mech.*, **75**, 1981, 27–38 (in Russian).

[17] Nenov, N. On the Zykov numbers and some its applications to Ramsey theory. *Serdica Bulg. Math. Publ.*, **9**, 1983, 161–167 (in Russian).

[18] Nenov, N. The chromatic number of any 10-vertex graph without 4-cliques is at most 4. *Compt. rend. Acad. bulg. Sci.*, **37**, 1984, 301–304 (in Russian).

[19] Nenov, N. On the small graphs with chromatic number 5 without 4-cliques. *Discrete Math.*, **188**, 1998, 297–298.

[20] Nenov, N. On a class of vertex Folkman graphs. *Annuaire Univ. Sofia Fac. Math. Inform.*, **94**, 2000, 15–25.

[21] Nenov, N. A generalization of a result of Dirac. *Ann. Univ. Sofia Fac. Math. Inform.*, **95**, 2001, 59–69.

[22] Nenov, N. On the 3-coloring vertex Folkman number $F(2, 2, 4)$. *Serdica Math. J.*, **27**, 2001, 131–136.

[23] Nenov, N. On a class of vertex Folkman numbers. *Serdica Math. J.*, **28**, 2002, 219–232.

[24] Nenov, N. On the triangle vertex Folkman numbers. *Discrete mathematics*, **271**, 2003, 327–334.

[25] Radziszowski, S. Small Ramsey numbers. *The Electronic Journal of Combinatorics*, Dynamic Survay, version 11, August 1, 2006.

Faculty of Mathematics and Informatics
St. Kl. Ohridski University of Sofia
5, J. Bourchier Blvd.
BG-1164 Sofia, Bulgaria
e-mail: `nenov@fmi.uni-sofia.bg`